

## Linear Algebra

1. Compute the (complex) eigenvalues and the associated (complex) eigenvectors for the following matrices:

$$\begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}, \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}, \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix} \text{ and } \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}.$$

2. The rank  $r(\mathbf{A})$  of a matrix  $\mathbf{A}$  is the maximal number of linearly independent column vectors in  $\mathbf{A}$ . Determine the rank of the following matrix for all values of the real parameter  $a$ . Hint: Use elementary row operations.

$$\mathbf{A} = \begin{pmatrix} 1 & -1 & 2 & 0 \\ -1 & a & -1 & a \\ 2 & 0 & 1 & -1 \end{pmatrix}$$

3. Verify the **Spectral Theorem for symmetric matrices** for the following matrix by finding a matrix  $\mathbf{P}$ :

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 0 \\ 1 & 1 & 0 \\ 0 & 0 & 2 \end{pmatrix}.$$

Investigate the definiteness of  $\mathbf{A}$ .

4. Use the identity  $\det(\mathbf{A} \cdot \mathbf{B}) = \det(\mathbf{A}) \cdot \det(\mathbf{B})$  for all  $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{n \times n}$  to prove the following facts:

- (a)  $\det(\mathbf{A}^{-1}) = (\det(\mathbf{A}))^{-1}$ ;
- (b)  $\det(\lambda \mathbf{A}) = \lambda^n \det(\mathbf{A})$  for  $\lambda \in \mathbb{R}$ ;
- (c)  $\mathbf{A}$  and  $\mathbf{P}^{-1} \mathbf{A} \mathbf{P}$  have the same eigenvalues.

5. Let  $\mathbf{A}$  be a square matrix with  $\det \mathbf{A} \neq 0$  and let  $\lambda$  be an eigenvalue of  $\mathbf{A}$ . Prove:

- (a)  $\lambda \neq 0$  and
- (b)  $1/\lambda$  is an eigenvalue of the inverse  $\mathbf{A}^{-1}$ .