

# Linear algebra

**Keywords:** vector, matrix, eigenvalue, eigenvector, diagonalization, linear transformation, quadratic forms and symmetric matrices

**Compare:** Vorlesung Mathematik 2

## Contents

<b>1</b>	<b>Matrices and vectors</b>	<b>2</b>
1.1	Real Vectors . . . . .	2
1.2	Real Matrices . . . . .	2
1.3	Linear transformations and matrices . . . . .	3
1.4	Complex matrices and vectors . . . . .	4
1.5	Matrix calculus . . . . .	4
<b>2</b>	<b>Eigenvalues and eigenvectors</b>	<b>5</b>
2.1	Definition and determination . . . . .	5
2.2	*Generalized Eigenvectors* . . . . .	7
<b>3</b>	<b>Diagonalization</b>	<b>9</b>
<b>4</b>	<b>Quadratic forms and matrices</b>	<b>14</b>

# 1 Matrices and vectors

## 1.1 Real Vectors

- $n$ -dimensional space  $\mathbb{R}^n$
- elements  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$  are called  $n$ -vectors

$$\mathbf{x} = \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = (x_1 \ x_2 \ \dots \ x_n)^T \text{ and } \mathbf{y} = \begin{pmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{pmatrix}$$

- scalar product and norm:

$$\begin{aligned} \mathbf{x} \bullet \mathbf{y} &= \langle \mathbf{x}, \mathbf{y} \rangle = x_1y_1 + x_2y_2 + \dots + x_ny_n \\ \|\mathbf{x}\| &= \sqrt{x_1^2 + x_2^2 + \dots + x_n^2} \end{aligned}$$

$$\mathbf{x} \bullet \mathbf{y} = \|\mathbf{x}\| \cdot \|\mathbf{y}\| \cdot \cos \angle(\mathbf{x}, \mathbf{y})$$

- $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k \in \mathbb{R}^n$

- If  $a_1, a_2, \dots, a_k \in \mathbb{R}$ , then  $\mathbf{z} = a_1\mathbf{x}_1 + a_2\mathbf{x}_2 + \dots + a_k\mathbf{x}_k$  is called a linear combination of  $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$ .
- $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$  are called linearly dependent, if there exist  $b_1, b_2, \dots, b_k \in \mathbb{R}$  such that  $b_1\mathbf{x}_1 + b_2\mathbf{x}_2 + \dots + b_k\mathbf{x}_k = \mathbf{0}$  and not all  $b_j = 0$ .
- $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_k$  are called linearly independent, if a linear combination of the zero vector

$$b_1\mathbf{x}_1 + b_2\mathbf{x}_2 + \dots + b_k\mathbf{x}_k = \mathbf{0}$$

is possible only with  $b_1 = b_2 = \dots = b_k = 0$ .

## 1.2 Real Matrices

$$\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_m \in \mathbb{R}^n$$

$$\mathbf{a}_1 = \begin{pmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{n1} \end{pmatrix}, \mathbf{a}_2 = \begin{pmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{n2} \end{pmatrix}, \dots, \mathbf{a}_m = \begin{pmatrix} a_{1m} \\ a_{2m} \\ \vdots \\ a_{nm} \end{pmatrix} \rightarrow \mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22} & \dots & a_{2m} \\ \vdots & \vdots & \vdots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nm} \end{pmatrix}$$

is called an  $n \times m$  matrix.

Notation:  $\mathbf{A} \in \mathbb{R}^{n \times m}$

- The inverse matrix  $\mathbf{A}^{-1}$  of the  $n \times n$  matrix  $\mathbf{A} = (a_{ij})$  is defined by

$$\mathbf{A}^{-1} \cdot \mathbf{A} = \mathbf{A} \cdot \mathbf{A}^{-1} = \mathbf{I}_n = \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{pmatrix}.$$

- For the  $n \times n$  matrix  $\mathbf{A}$  let  $\mathbf{A}_{ij}$  denote the  $(n-1) \times (n-1)$  submatrix of  $\mathbf{A}$  generated by cancelling the  $i$ -th row and the  $j$ -th column of  $\mathbf{A}$ . Then the determinant  $\det(\mathbf{A})$  is given (recursively) by

$$\det(\mathbf{A}) = |\mathbf{A}| = a_{11} \det \mathbf{A}_{11} - a_{12} \det \mathbf{A}_{12} + \cdots + (-1)^{n+1} a_{1n} \det \mathbf{A}_{1n}$$

- $\det(\mathbf{A} \cdot \mathbf{B}) = \det(\mathbf{A}) \cdot \det(\mathbf{B})$

### Example 1.1

$$\begin{aligned} & \left| \begin{array}{cccc} 1 & 1 & 3 & 3 \\ 1 & 2 & 1 & 2 \\ 1 & -2 & 1 & -2 \\ 0 & 1 & -2 & -1 \end{array} \right| \\ &= 1 \cdot \left| \begin{array}{ccc} 2 & 1 & 2 \\ -2 & 1 & -2 \\ 1 & -2 & -1 \end{array} \right| - 1 \cdot \left| \begin{array}{ccc} 1 & 1 & 2 \\ 1 & 1 & -2 \\ 0 & -2 & -1 \end{array} \right| + 3 \cdot \left| \begin{array}{ccc} 1 & 2 & 2 \\ 1 & -2 & -2 \\ 0 & 1 & -1 \end{array} \right| - 3 \cdot \left| \begin{array}{ccc} 1 & 2 & 1 \\ 1 & -2 & 1 \\ 0 & 1 & -2 \end{array} \right|. \end{aligned}$$

## 1.3 Linear transformations and matrices

**Definition 1.1** A linear transformation is a map  $T : \mathbb{R}^m \rightarrow \mathbb{R}^n$  such that for all  $\mathbf{x}, \mathbf{y} \in \mathbb{R}^m$  and all  $\lambda, \mu \in \mathbb{R}$  we have:

$$T(\lambda \cdot \mathbf{x} + \mu \cdot \mathbf{y}) = \lambda \cdot T(\mathbf{x}) + \mu \cdot T(\mathbf{y})$$

Each  $n \times m$  matrix  $\mathbf{A}$  defines a linear transformation by matrix multiplication

$$T_{\mathbf{A}}(\mathbf{x}) = \mathbf{A} \cdot \mathbf{x} = x_1 \mathbf{a}_1 + \cdots + x_m \mathbf{a}_m.$$

The image of the vector  $\mathbf{x} \in \mathbb{R}^m$  is a linear combination of the column vectors of the matrix  $\mathbf{A}$ .

## 1.4 Complex matrices and vectors

Sometimes it is helpful to allow complex matrices and vectors (matrices whose elements are complex numbers). A complex matrix can be viewed as a combination of two real matrices:

$$\begin{aligned} \mathbf{A} &= \begin{pmatrix} a_{11} + ib_{11} & a_{12} + ib_{12} & \dots & a_{1m} + ib_{1m} \\ a_{21} + ib_{21} & a_{22} + ib_{22} & \dots & a_{2m} + ib_{2m} \\ \vdots & \vdots & \vdots & \vdots \\ a_{n1} + ib_{n1} & a_{n2} + ib_{n2} & \dots & a_{nm} + ib_{nm} \end{pmatrix} \\ &= \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1m} \\ a_{21} & a_{22} & \dots & a_{2m} \\ \vdots & \vdots & \vdots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nm} \end{pmatrix} + i \cdot \begin{pmatrix} b_{11} & b_{12} & \dots & b_{1m} \\ b_{21} & b_{22} & \dots & b_{2m} \\ \vdots & \vdots & \vdots & \vdots \\ b_{n1} & b_{n2} & \dots & b_{nm} \end{pmatrix} \end{aligned}$$

## 1.5 Matrix calculus

- 1a.  $\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$
- 1b. In general:  $\mathbf{AB} \neq \mathbf{BA}$
- 2a.  $(\mathbf{A} + \mathbf{B}) + \mathbf{C} = \mathbf{A} + (\mathbf{B} + \mathbf{C})$
- 2b.  $(\mathbf{AB})\mathbf{C} = \mathbf{A}(\mathbf{BC})$
- 3a.  $\mathbf{A} + \mathbf{0} = \mathbf{A}$
- 3b.  $\mathbf{AI} = I\mathbf{A} = \mathbf{A}$  (  $\mathbf{A}$  square )
- 4.  $\mathbf{AB} = \mathbf{0} \nRightarrow \mathbf{A} = \mathbf{0}$  or  $\mathbf{B} = \mathbf{0}$
- 5.  $\mathbf{AB} = \mathbf{AC} \nRightarrow \mathbf{B} = \mathbf{C}$
- 6.  $\lambda(\mathbf{A} + \mathbf{B}) = \lambda\mathbf{A} + \lambda\mathbf{B} \quad \lambda \in \mathbb{R}$
- 7.  $\mathbf{A}(\mathbf{B} + \mathbf{C}) = \mathbf{AB} + \mathbf{AC}$
- 8.  $(\mathbf{A} + \mathbf{B})\mathbf{C} = \mathbf{AC} + \mathbf{BC}$
- 9.  $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$
- 10.  $(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$
- 11.  $(\mathbf{A}^T)^T = \mathbf{A}$
- 12.  $(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T$
- 13.  $(\mathbf{AB})^T = \mathbf{B}^T\mathbf{A}^T$
- 14.  $(\mathbf{A}^{-1})^T = (\mathbf{A}^T)^{-1}$

For  $\mathbf{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  with  $ad - bc \neq 0$  is  $\mathbf{A}^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}$ .

All these definitions and results can be generalized to vectors and matrices with complex entries.

## 2 Eigenvalues and eigenvectors

### 2.1 Definition and determination

**Definition 2.1** If  $\mathbf{A}$  is a real (or complex)  $n \times n$  matrix, then a (complex) number  $\lambda$  is an eigenvalue of  $\mathbf{A}$  if there is a nonzero (complex) vector  $\mathbf{x} \in \mathbb{C}^n$  such that

$$\boxed{\mathbf{A}\mathbf{x} = \lambda\mathbf{x}}$$

Then  $\mathbf{x}$  is an eigenvector of  $\mathbf{A}$  (associated with  $\lambda$ ).

Remark: If  $\mathbf{x}$  is an eigenvector associated with the eigenvalue  $\lambda$ , then so is  $\alpha\mathbf{x}$  for every real (and complex) number  $\alpha \neq 0$ .

$$\mathbf{A}(\alpha\mathbf{x}) = \alpha\mathbf{A}\mathbf{x} = \alpha(\lambda\mathbf{x}) = \lambda(\alpha\mathbf{x})$$

How to find eigenvalues? The equation can be written as

$$\begin{aligned} \mathbf{A}\mathbf{x} &= \lambda\mathbf{x} \\ \Leftrightarrow \mathbf{A}\mathbf{x} - \lambda\mathbf{I}\mathbf{x} &= \mathbf{0} \\ \Leftrightarrow (\mathbf{A} - \lambda\mathbf{I})\mathbf{x} &= \mathbf{0} \end{aligned}$$

This is a homogeneous system of linear equations. It has a solution  $\mathbf{x} \neq \mathbf{0}$  if and only if the matrix  $(\mathbf{A} - \lambda\mathbf{I})$  is singular which means that its determinant equals to 0.

$$\boxed{(\mathbf{A} - \lambda\mathbf{I}) \text{ singular} \Leftrightarrow \underbrace{\det(\mathbf{A} - \lambda\mathbf{I})}_{p_A(\lambda)} = 0}$$

$p_A(\lambda) = 0$  is called the characteristic equation of  $\mathbf{A}$ . The function  $p_A(\lambda)$  is a polynomial of degree  $n$  in  $\lambda$ , called the characteristic polynomial of  $\mathbf{A}$ .

#### Determination of the eigenvalues and eigenvectors

1. The polynomial equation  $p_A(\lambda) = 0$  has always  $n$  complex solutions (counted with multiplicity) and may have no real solutions. If  $\lambda_1, \dots, \lambda_r \in \mathbb{C}$  are the pairwise distinct solutions (the eigenvalues of  $\mathbf{A}$ ) with the multiplicities  $k_1, \dots, k_r$  then the characteristic polynomial can be written as

$$p_A(\lambda) = (\lambda_1 - \lambda)^{k_1} (\lambda_2 - \lambda)^{k_2} \dots (\lambda_r - \lambda)^{k_r}.$$

The multiplicity  $k_i$  of the zero  $\lambda_i$  is called algebraic multiplicity of the eigenvalue  $\lambda_i$ . Generally, the determination of the (exact) zeros is impossible for  $n \geq 5$  and we have to use numerical methods.

2. For each eigenvalue  $\lambda_i$  ( $1 \leq i \leq r$ ) we compute the so-called eigenspace for  $\lambda_i$

$$V(\lambda_i) = \{ \mathbf{x} \in \mathbb{C}^n \mid (\mathbf{A} - \lambda_i \mathbf{I}) \mathbf{x} = \mathbf{0} \}.$$

The dimension of the vector space  $V(\lambda_i)$  is called the geometric multiplicity of the eigenvalue  $\lambda_i$ .

**Example 2.1**

$$A = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -6 & -1 & 4 \end{pmatrix}$$

- $p_A(\lambda) = -\lambda^3 + 4\lambda^2 - \lambda - 6 = (\lambda + 1) \cdot (-\lambda^2 + 5\lambda - 6) = -(\lambda + 1) \cdot (\lambda - 2) \cdot (\lambda - 3)$
- *Zeros of the characteristic polynomial:  $\lambda_1 = -1$ ,  $\lambda_2 = 2$  and  $\lambda_3 = 3$  (all of algebraic multiplicity 1)*

•

$$\left[ \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -6 & -1 & 4 \end{pmatrix} - (-1) \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right] \cdot \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \mathbf{x}^{(1)} = \begin{pmatrix} 1 \\ -1 \\ 1 \end{pmatrix}$$

and  $V(-1) = \{ t \cdot \mathbf{x}^{(1)} \mid t \in \mathbb{R} \}$  with geometric multiplicity 1.

•

$$\left[ \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -6 & -1 & 4 \end{pmatrix} - 2 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right] \cdot \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \mathbf{x}^{(2)} = \begin{pmatrix} 1 \\ 2 \\ 4 \end{pmatrix}$$

and  $V(2) = \{ t \cdot \mathbf{x}^{(2)} \mid t \in \mathbb{R} \}$  with geometric multiplicity 1.

•

$$\left[ \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ -6 & -1 & 4 \end{pmatrix} - 3 \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right] \cdot \begin{pmatrix} x_1 \\ x_2 \\ x_3 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} \rightarrow \mathbf{x}^{(3)} = \begin{pmatrix} 1 \\ 3 \\ 9 \end{pmatrix}$$

and  $V(3) = \{ t \cdot \mathbf{x}^{(3)} \mid t \in \mathbb{R} \}$  with geometric multiplicity 1.

**Definition 2.2** *The spectral radius of a quadratic matrix  $A$  is the real number*

$$\rho(A) := \max\{|\lambda_1|, \dots, |\lambda_r|\}.$$

## 2.2 \*Generalized Eigenvectors\*

To solve some interesting problems we have to generalize the notion of eigenvectors.

**Definition 2.3** A vector  $\mathbf{x} \in \mathbb{C}^n$  is called generalized eigenvector of degree  $l \in \mathbb{N}$  associated to the eigenvalue  $\lambda$  of  $\mathbf{A}$ , if

$$(\mathbf{A} - \lambda \mathbf{I})^l \mathbf{x} = \mathbf{0} \quad \text{and} \quad (\mathbf{A} - \lambda \mathbf{I})^{l-1} \mathbf{x} \neq \mathbf{0}.$$

Of course, an eigenvector is a generalized eigenvector of degree 1.

**Example 2.2** The matrix

$$\mathbf{A} = \begin{pmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}$$

has the eigenvalue 1 of (algebraic) multiplicity 3 with  $\dim V(1) = 1$  (geometric multiplicity). We have:

$$\begin{aligned} (\mathbf{A} - \mathbf{I}) \mathbf{e}_1 &= \mathbf{0} & (\mathbf{A} - \mathbf{I}) \mathbf{e}_2 &= \mathbf{e}_1 & (\mathbf{A} - \mathbf{I})^2 \mathbf{e}_2 &= \mathbf{0} \\ (\mathbf{A} - \mathbf{I}) \mathbf{e}_3 &= \mathbf{e}_1 + \mathbf{e}_2 & (\mathbf{A} - \mathbf{I})^2 \mathbf{e}_3 &= \mathbf{e}_1 & (\mathbf{A} - \mathbf{I})^3 \mathbf{e}_3 &= \mathbf{0} \end{aligned}$$

This means, that  $\mathbf{e}_1$  is an eigenvector,  $\mathbf{e}_2$  is a generalized eigenvector of degree 2 and  $\mathbf{e}_3$  is a generalized eigenvector of degree 3.

**Theorem 2.1** Let  $\mathbf{A} \in \mathbb{C}^{n \times n}$  be a complex (or real) matrix with

$$p_A(\lambda) = (\lambda_1 - \lambda)^{k_1} (\lambda_2 - \lambda)^{k_2} \dots (\lambda_r - \lambda)^{k_r}.$$

- Let  $\lambda$  be an eigenvalue of  $\mathbf{A}$  of (algebraic) multiplicity  $l$ . Then there exist  $l$  linearly independent generalized eigenvectors (of degree  $\leq l$ ). This means:

$$\dim \{ \mathbf{x} \in \mathbb{C}^n \mid (\mathbf{A} - \lambda \mathbf{I})^l \mathbf{x} = \mathbf{0} \} = l.$$

- Generalized eigenvectors associated to pairwise different eigenvalues of  $\mathbf{A}$  are linearly independent.
- There exists a basis  $\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_n$  of  $\mathbb{C}^n$  consisting of generalized eigenvectors of  $\mathbf{A}$ . If  $\mathbf{P}$  is the matrix with this basis as the columns, then

$$\mathbf{P}^{-1} \mathbf{A} \mathbf{P} = \begin{pmatrix} \boxed{\mathbf{A}_1} & & & \mathbf{0} \\ & \boxed{\mathbf{A}_2} & & \\ & & \ddots & \\ \mathbf{0} & & & \boxed{\mathbf{A}_r} \end{pmatrix}$$

with  $\mathbf{A}_i \in \mathbb{C}^{k_i \times k_i}$  for all  $i = 1, 2, \dots, r$ .

Let us have a look at the case  $n = 2$  and  $\mathbf{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ .

1. Characteristic polynomial:

$$\begin{aligned} p_A(\lambda) &= \det \begin{pmatrix} a - \lambda & b \\ c & d - \lambda \end{pmatrix} \\ &= \underbrace{\lambda^2 - (a + d)\lambda + ad - bc}_{=:tr(A)} = \underbrace{(a + d)\lambda + ad - bc}_{=\det(A)} = (\lambda_1 - \lambda)(\lambda_2 - \lambda) \end{aligned}$$

$$\text{with } \lambda_{1,2} = \frac{a + d}{2} \pm \sqrt{\frac{(a + d)^2}{4} - \det(A)}.$$

2. For each  $\lambda_i$  ( $i = 1, 2$ ) we solve the linear system

$$\begin{pmatrix} a - \lambda_i & b \\ c & d - \lambda_i \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

We have four different cases:

1.  $\lambda_1, \lambda_2 \in \mathbb{R}, \lambda_1 \neq \lambda_2$

$$\text{Example: } \mathbf{A} = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$$

We have  $p_A(\lambda) = (1 - \lambda)^2 - 4 = (\lambda + 1)(\lambda - 3)$  (two different eigenvalues of algebraic multiplicity 1). A direct calculation shows, that  $\dim V(-1) = 1$  and  $\dim V(3) = 1$  and the geometric multiplicity are (of all eigenvalues) equal to the algebraic multiplicity.

2.  $\lambda = \lambda_1 = \lambda_2 \in \mathbb{R}$  with  $\dim V(\lambda) = 2$

$$\text{Example: } \mathbf{A} = \begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$$

We have  $p_A(\lambda) = (2 - \lambda)^2$  (one eigenvalue of algebraic multiplicity 2). A direct calculation shows, that  $\dim V(2) = 2$  and the geometric multiplicity (of the eigenvalue 2) is equal to the algebraic multiplicity.

3.  $\lambda = \lambda_1 = \lambda_2 \in \mathbb{R}$  with  $\dim V(\lambda) = 1$

$$\text{Example: } \mathbf{A} = \begin{pmatrix} 2 & 1 \\ 0 & 2 \end{pmatrix}$$

We have  $p_A(\lambda) = (2 - \lambda)^2$  (one eigenvalue of algebraic multiplicity 2). A direct calculation shows, that  $\dim V(2) = 1$  and the geometric multiplicity of the eigenvalue 2 is different of the algebraic multiplicity.

4.  $\lambda_2 = \overline{\lambda_1} \in \mathbb{C} - \mathbb{R}$

$$\text{Example: } \mathbf{A} = \begin{pmatrix} \cos \phi & -\sin \phi \\ \sin \phi & \cos \phi \end{pmatrix} \text{ with } \phi \neq k\pi$$

We have  $p_A(\lambda) = (\lambda - \cos \phi)^2 + \sin^2 \phi = \lambda^2 - 2\lambda \cos \phi + 1$  with the two different complex zeroes  $\lambda_{1,2} = \cos \phi \pm i \sin \phi$ .

### 3 Diagonalization

Let  $\mathbf{A}$  and  $\mathbf{P}$  be  $n \times n$  matrices with  $\mathbf{P}$  invertible. Then  $\mathbf{A}$  and  $\mathbf{P}^{-1}\mathbf{A}\mathbf{P}$  have the same eigenvalues (because they have the same characteristic polynomial).

**Definition 3.1** An  $n \times n$  matrix  $\mathbf{A}$  is diagonalizable if there is an invertible matrix  $\mathbf{P}$  and a diagonal matrix  $\mathbf{D}$  such that

$$\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \mathbf{D}.$$

Two natural questions:

1. Which square matrices are diagonalizable?
2. If  $\mathbf{A}$  is diagonalizable, how do we find the matrix  $\mathbf{P}$ ?

**Theorem 3.1** An  $n \times n$  matrix  $\mathbf{A}$  is diagonalizable if and only if it has a set of  $n$  linearly independent eigenvectors  $\mathbf{p}_1, \dots, \mathbf{p}_n$ . In this case,

$$\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{pmatrix},$$

where  $\mathbf{P}$  is the matrix with  $\mathbf{p}_1, \dots, \mathbf{p}_n$  as its columns, and  $\lambda_1, \dots, \lambda_n$  are the corresponding eigenvalues.

**Proof:** We prove only one direction of the statement:

$\mathbf{A}$  has  $n$  linearly independent eigenvectors  $\implies \mathbf{A}$  is diagonalizable.

Let  $\mathbf{p}_1, \mathbf{p}_2, \dots, \mathbf{p}_n$  be the  $n$  linearly independent eigenvectors of  $\mathbf{A}$  with corresponding eigenvalues  $\lambda_1, \lambda_2, \dots, \lambda_n$ . We form the matrix

$$\mathbf{P} = \begin{pmatrix} | & | & & | \\ \mathbf{p}_1 & \mathbf{p}_2 & \dots & \mathbf{p}_n \\ | & | & & | \end{pmatrix}$$

with the eigenvectors of  $\mathbf{A}$  as the columns. Then

$$\mathbf{AP} = \begin{pmatrix} | & | & & | \\ \mathbf{Ap}_1 & \mathbf{Ap}_2 & \dots & \mathbf{Ap}_n \\ | & | & & | \end{pmatrix}$$

the column vectors of  $\mathbf{AP}$  are the vectors  $\mathbf{Ap}_1, \mathbf{Ap}_2, \dots, \mathbf{Ap}_n$ . Using the

property of eigenvectors, we get

$$\begin{aligned}
 \mathbf{AP} &= \begin{pmatrix} | & | & | & | \\ \mathbf{Ap}_1 & \mathbf{Ap}_2 & \dots & \mathbf{Ap}_n \\ | & | & & | \end{pmatrix} \\
 &= \begin{pmatrix} | & | & | & | \\ \lambda_1 \mathbf{p}_1 & \lambda_2 \mathbf{p}_2 & \dots & \lambda_n \mathbf{p}_n \\ | & | & & | \end{pmatrix} \\
 &= \begin{pmatrix} | & | & | & | \\ \mathbf{p}_1 & \mathbf{p}_2 & \dots & \mathbf{p}_n \\ | & | & & | \end{pmatrix} \begin{pmatrix} \lambda_1 & & & 0 \\ & \lambda_2 & & \\ & & \ddots & \\ 0 & & & \lambda_n \end{pmatrix} \\
 &= \mathbf{PD}.
 \end{aligned}$$

where  $\mathbf{D}$  is the diagonal matrix with diagonal entries equal to the eigenvalues of  $\mathbf{A}$ . The matrix  $\mathbf{P}$  has maximal rank (and is invertible), because the column vectors are linearly independent. Hence the equation  $\mathbf{AP} = \mathbf{PD}$  is equivalent to  $\mathbf{P}^{-1}\mathbf{AP} = \mathbf{D}$ .

□

**Example 3.1** The matrix  $\mathbf{A} = \begin{pmatrix} 1 & 1 \\ -2 & 4 \end{pmatrix}$  has the eigenvalues and eigenvectors

$$\begin{aligned}
 \lambda_1 &= 2 & \mathbf{p}_1 &= \begin{pmatrix} 1 \\ 1 \end{pmatrix} \\
 \lambda_2 &= 3 & \mathbf{p}_2 &= \begin{pmatrix} 1 \\ 2 \end{pmatrix}
 \end{aligned}$$

Hence  $\mathbf{P} = \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ ,  $\mathbf{P}^{-1} = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix}$  and:

$$\mathbf{P}^{-1}\mathbf{AP} = \begin{pmatrix} 2 & -1 \\ -1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ -2 & 4 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 2 & 0 \\ 0 & 3 \end{pmatrix}$$

Many matrices encountered in economics are (real) symmetric and for these matrices we have the following important result.

**Theorem 3.2 (Spectral Theorem for symmetric matrices)** *If the real  $n \times n$  matrix  $\mathbf{A}$  is symmetric ( $\mathbf{A} = \mathbf{A}^T$ ), then:*

1. *All  $n$  eigenvalues  $\lambda_1, \dots, \lambda_n$  are real.*
2. *Eigenvectors that correspond to different eigenvalues are orthogonal.*
3. *There exists an orthogonal and real matrix  $\mathbf{P}$  ( $\mathbf{P}^{-1} = \mathbf{P}^T$ ) such that*

$$\mathbf{P}^{-1} \mathbf{A} \mathbf{P} = \begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{pmatrix}.$$

The columns  $\mathbf{p}_1, \dots, \mathbf{p}_n$  of the matrix  $\mathbf{P}$  are eigenvectors of unit length corresponding to the eigenvalues  $\lambda_1, \dots, \lambda_n$ .

**Proof:** Let  $\mathbf{A}$  be a real and symmetric  $n \times n$  matrix.

1. Let  $\mathbf{A}\mathbf{p}_i = \lambda_i \mathbf{p}_i$ . By complex conjugation of this equation (complex conjugate all entries of the vector and matrix, but keep in mind that  $\mathbf{A}$  has only real entries) we get

$$\overline{\mathbf{A}\mathbf{p}_i} = \overline{\mathbf{A}\overline{\mathbf{p}_i}} = \mathbf{A}\overline{\mathbf{p}_i} = \overline{\lambda_i \mathbf{p}_i}$$

and

$$\lambda_i \mathbf{p}_i^T \overline{\mathbf{p}_i} = (\mathbf{A}\mathbf{p}_i)^T \overline{\mathbf{p}_i} = \mathbf{p}_i^T \mathbf{A}^T \overline{\mathbf{p}_i} = \mathbf{p}_i^T \mathbf{A} \overline{\mathbf{p}_i} = \mathbf{p}_i^T \overline{\lambda_i \mathbf{p}_i} = \overline{\lambda_i} \mathbf{p}_i^T \overline{\mathbf{p}_i}$$

Because  $\mathbf{p}_i^T \overline{\mathbf{p}_i} = \|\mathbf{p}_i\|^2 \neq 0$ , we have  $\lambda_i = \overline{\lambda_i}$  and  $\lambda_i$  must be a real number.

2. Let  $\mathbf{A}\mathbf{p}_i = \lambda_i \mathbf{p}_i$  and  $\mathbf{A}\mathbf{p}_j = \lambda_j \mathbf{p}_j$  with  $\lambda_i \neq \lambda_j$ . Then

$$\begin{aligned} \lambda_i \mathbf{p}_i^T \mathbf{p}_j &= (\mathbf{A}\mathbf{p}_i)^T \mathbf{p}_j \\ &= \mathbf{p}_i^T \mathbf{A}^T \mathbf{p}_j \\ &= \mathbf{p}_i^T (\mathbf{A}^T \mathbf{p}_j) \\ &= \mathbf{p}_i^T (\mathbf{A}\mathbf{p}_j) \quad \text{because } \mathbf{A} = \mathbf{A}^T \\ &= \mathbf{p}_i^T \lambda_j \mathbf{p}_j \\ &= \lambda_j \mathbf{p}_i^T \mathbf{p}_j \end{aligned}$$

or

$$\lambda_i (\mathbf{p}_i^T \mathbf{p}_j) = \lambda_j (\mathbf{p}_i^T \mathbf{p}_j)$$

and because  $\lambda_i \neq \lambda_j$ , the scalar product of  $\mathbf{p}_i$  and  $\mathbf{p}_j$  must be zero:  $\mathbf{p}_i^T \mathbf{p}_j = \mathbf{p}_i \bullet \mathbf{p}_j = 0$ . Hence the two eigenvectors are orthogonal.

3. We give the proof of part 3 only for the case that all eigenvalues  $\lambda_1, \dots, \lambda_n$  are (pairwise) different (and real by part 1). In this case, the corresponding eigenvectors  $\mathbf{p}'_1, \dots, \mathbf{p}'_n$  are orthogonal (by part 2) and hence linearly independent. Now choose for  $i = 1, \dots, n$  an eigenvector of length 1 by

$$\mathbf{p}_i := \frac{1}{\|\mathbf{p}'_i\|} \mathbf{p}'_i$$

It is easy to show, that

$$\mathbf{p}_i^T \mathbf{p}_j = \mathbf{p}_i \bullet \mathbf{p}_j = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

The matrix

$$\mathbf{P} = \begin{pmatrix} | & | & & | \\ \mathbf{p}_1 & \mathbf{p}_2 & \cdots & \mathbf{p}_n \\ | & | & & | \end{pmatrix}$$

is an orthogonal matrix, because

$$\begin{aligned} \mathbf{P}^T \mathbf{P} &= \begin{pmatrix} - & \mathbf{p}_1^T & - \\ - & \mathbf{p}_2^T & - \\ \dots & & \\ - & \mathbf{p}_n^T & - \end{pmatrix} \begin{pmatrix} | & | & & | \\ \mathbf{p}_1 & \mathbf{p}_2 & \cdots & \mathbf{p}_n \\ | & | & & | \end{pmatrix} \\ &= \begin{pmatrix} \mathbf{p}_1^T \mathbf{p}_1 & \mathbf{p}_1^T \mathbf{p}_2 & \cdots & \mathbf{p}_1^T \mathbf{p}_n \\ \mathbf{p}_2^T \mathbf{p}_1 & \mathbf{p}_2^T \mathbf{p}_2 & \cdots & \mathbf{p}_2^T \mathbf{p}_n \\ \dots & \dots & \ddots & \dots \\ \mathbf{p}_n^T \mathbf{p}_1 & \mathbf{p}_n^T \mathbf{p}_2 & \cdots & \mathbf{p}_n^T \mathbf{p}_n \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \dots & \dots & \ddots & \dots \\ 0 & 0 & \cdots & 1 \end{pmatrix}. \end{aligned}$$

Hence we have  $\mathbf{P}^T = \mathbf{P}^{-1}$

□

**Example 3.2** The matrix  $\mathbf{A} = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$  is symmetric and has the eigenvalues  $\lambda_1 = -1$  and  $\lambda_2 = 3$ . The corresponding eigenspaces are

$$V(-1) = t_1 \begin{pmatrix} 1 \\ -1 \end{pmatrix} \quad V(3) = t_2 \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

The two eigenspaces are orthogonal, because the scalar product of the two spanning vectors is 0. In order to construct the matrix  $\mathbf{P}$ , we have to use eigenvectors of length 1 (unit vectors). A spanning vector of length 1 for  $V(-1)$  is

$$\mathbf{p}_1 = \frac{1}{\sqrt{1^2 + (-1)^2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ -1 \end{pmatrix}$$

and for  $V(3)$  is

$$\mathbf{p}_2 = \frac{1}{\sqrt{1^2 + 1^2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

Hence  $\mathbf{P} = \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ -1/\sqrt{2} & 1/\sqrt{2} \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}$  is an orthogonal matrix, because

$\mathbf{P}^{-1} = \mathbf{P}^T = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix}$  and

$$\mathbf{P}^{-1} \mathbf{A} \mathbf{P} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix} \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix} = \begin{pmatrix} -1 & 0 \\ 0 & 3 \end{pmatrix}$$

## 4 Quadratic forms and matrices

**Definition 4.1** A quadratic form in  $n$  variables  $\mathbf{x} = (x_1, \dots, x_n)^T$  is a function of the form

$$Q_{\mathbf{A}}(\mathbf{x}) = \sum_{i,j=1}^n a_{ij}x_i x_j = \mathbf{x}^T \mathbf{A} \mathbf{x}$$

where  $\mathbf{A} = (a_{ij})$  is an  $n \times n$  matrix.

Quadratic forms are important examples of multi-variate functions and  $Q_{\mathbf{A}}$  is a homogeneous function of degree 2 in  $n$  variables.

**Example 4.1**  $Q(x_1, x_2) = x_1^2 + x_1 x_2 + x_2^2$  is a quadratic form and can be written as

$$\begin{aligned} (x_1 \ x_2) \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} &= (x_1 \ x_2) \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \\ &= (x_1 \ x_2) \begin{pmatrix} 1 & 1/2 \\ 1/2 & 1 \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} \\ &= \dots \end{aligned}$$

Unfortunaetly, there is no unique way to write a given quadratic form in matrix term. But we may resolve this situation by **always choosing  $\mathbf{A}$  to be symmetric!**


**Exercise 4.1** Let  $Q(\mathbf{x}) = \mathbf{x}^T \mathbf{B} \mathbf{x}$  where  $\mathbf{B}$  is not symmetric. Let  $\mathbf{A} = (\mathbf{B} + \mathbf{B}^T)/2$  and  $\mathbf{C} = (\mathbf{B} - \mathbf{B}^T)/2$ . Show that  $\mathbf{A}$  is symmetric and evaluate both  $\mathbf{x}^T \mathbf{A} \mathbf{x}$  and  $\mathbf{x}^T \mathbf{C} \mathbf{x}$ .

### Example 4.2


- The quadratic form  $Q(x_1, x_2) = x_1^2 + x_1x_2 + x_2^2$  can be written as

$$\left(x_1 + \frac{x_2}{2}\right)^2 + \frac{3}{4}x_2^2.$$


As a sum of squares, it can not be negative and can only be zero when  $x_1 + \frac{x_2}{2} = 0$  and  $x_2 = 0$ , or  $x_1 = x_2 = 0$ . We call this a positive definite quadratic form.



- The quadratic form  $Q(x_1, x_2) = x_1^2 + 2x_1x_2 + x_2^2 = (x_1 + x_2)^2$  is always non-negative, but it is zero whenever  $x_1 + x_2 = 0$  or  $x_1 = -x_2$  (it is zero for non-zero values of the variables). We call this a positive semi-definite quadratic form.



- The quadratic form  $Q(x_1, x_2) = x_1^2 - 6x_1x_2 = (x_1 - 3x_2)^2 - 9x_2^2$  can be positive or negative. We call this an indefinite quadratic form.



**Definition 4.2** A quadratic form  $Q_{\mathbf{A}}(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$ , as well as its associated symmetric matrix  $\mathbf{A}$ , is said to be

$$\begin{array}{ll} \text{positive definite} & : \iff Q_{\mathbf{A}}(\mathbf{x}) > 0 \\ \text{positive semi-definite} & : \iff Q_{\mathbf{A}}(\mathbf{x}) \geq 0 \\ \text{negative definite} & : \iff Q_{\mathbf{A}}(\mathbf{x}) < 0 \\ \text{negative semi-definite} & : \iff Q_{\mathbf{A}}(\mathbf{x}) \leq 0 \end{array}$$

for all  $\mathbf{x} \neq \mathbf{0}$ .

The quadratic form is called indefinite, if there are vectors  $\mathbf{a}$  and  $\mathbf{b}$  with  $Q_{\mathbf{A}}(\mathbf{a}) < 0$  and  $Q_{\mathbf{A}}(\mathbf{b}) > 0$ .

It is easy to see, that for  $i = 1, \dots, n$ :

$$Q_{\mathbf{A}}(\mathbf{e}_i) = a_{ii}.$$

The technique used in the examples to examine the sign of the quadratic form is known as **completing the squares**. Let us examine the possible signs of a quadratic form  $Q_{\mathbf{A}}(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$  using the eigenvalues/eigenvectors of the **symmetric** matrix  $\mathbf{A}$ .

By the **Spectral Theorem for symmetric matrices** we can choose a matrix  $\mathbf{P}$  of eigenvectors  $\mathbf{p}_1, \dots, \mathbf{p}_n$  of  $\mathbf{A}$ , such that  $\mathbf{P}^{-1} = \mathbf{P}^T$  and

$$\mathbf{P}^{-1} \mathbf{A} \mathbf{P} = \mathbf{P}^T \mathbf{A} \mathbf{P} = \begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{pmatrix},$$

where  $\lambda_1, \dots, \lambda_n$  are the eigenvalues of  $\mathbf{A}$ .

Now let  $\mathbf{y} := \mathbf{P}^T \mathbf{x}$ . This defines new variables  $y_1, \dots, y_n$  as linear combinations of the old ones

$$y_i = \sum_{j=1}^n p_{ji} x_j.$$

Further, since  $\mathbf{P} \mathbf{P}^T = \mathbf{I}$  we have  $\mathbf{x} = \mathbf{P} \mathbf{y}$  and

$$\begin{aligned} Q_{\mathbf{A}}(\mathbf{x}) &= \mathbf{x}^T \mathbf{A} \mathbf{x} \\ &= (\mathbf{P} \mathbf{y})^T \mathbf{A} (\mathbf{P} \mathbf{y}) \\ &= \mathbf{y}^T \mathbf{P}^T \mathbf{A} \mathbf{P} \mathbf{y} \\ &= \mathbf{y}^T (\mathbf{P}^T \mathbf{A} \mathbf{P}) \mathbf{y} \\ &= \mathbf{y}^T \begin{pmatrix} \lambda_1 & & 0 \\ & \ddots & \\ 0 & & \lambda_n \end{pmatrix} \mathbf{y} \\ &= \sum_{i=1}^n \lambda_i y_i^2. \end{aligned}$$

Thus we completed the squares. The quadratic form is expressed in terms of the new variables as a sum/difference of pure square terms. To determine the sign of the quadratic form, we simply inspect the signs of the eigenvalues of  $\mathbf{A}$ .

**Theorem 4.1 (Sylvester)**

If  $\mathbf{A}$  is symmetric, then the quadratic form  $Q_{\mathbf{A}}(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$  is

$$\begin{array}{ll} \text{positive definite} & \iff \forall \lambda_i > 0 \\ \text{positive semi-definite} & \iff \forall \lambda_i \geq 0 \\ \text{negative definite} & \iff \forall \lambda_i < 0 \\ \text{negative semi-definite} & \iff \forall \lambda_i \leq 0 \\ \text{indefinite} & \iff \exists \lambda_i > 0 \text{ and } \lambda_j < 0. \end{array}$$

Checking eigenvalues can be tedious. There is a convenient condition on the matrix  $\mathbf{A}$  in terms of certain sub-determinants, which can be used to identify the definiteness of  $\mathbf{A}$ .

An arbitrary principal minor of order  $r$  of an  $n \times n$  matrix  $\mathbf{A}$  is the determinant of a matrix obtained by deleting  $n - r$  rows and  $n - r$  columns of  $\mathbf{A}$  such that if the  $i$ th row (column) is selected then so is the  $i$ th column (row). A principal minor is called a leading principal minor of order  $r$  if it consists of the first (leading)  $r$  rows and columns of  $\mathbf{A}$ .

**Example 4.3** Let

$$\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$$

The principal minors of  $\mathbf{A}$  are  $\det(\mathbf{A})$ ,  $\det \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$ ,  $\det \begin{pmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{pmatrix}$ ,  $\det \begin{pmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{pmatrix}$ ,  $a_{11}$ ,  $a_{22}$  and  $a_{33}$ .

The leading principal minors are  $a_{11}$ ,  $\det \begin{pmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{pmatrix}$  and  $\det(\mathbf{A})$ .

**Theorem 4.2**

Let  $\mathbf{A}$  be a symmetric  $n \times n$  matrix. We denote by  $D_k$  the leading principal minor of order  $k$  and let  $\Delta_k$  denote an arbitrary principal minor of order  $k$ . Then the quadratic form  $Q_{\mathbf{A}}(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$  is

$$\begin{array}{ll} \text{positive definite} & \iff D_k > 0 \text{ for } k = 1, \dots, n \\ \text{positive semi-definite} & \iff \Delta_k \geq 0 \text{ for all principal minors of order } k = 1, \dots, n \\ \text{negative definite} & \iff (-1)^k D_k > 0 \text{ for } k = 1, \dots, n \\ \text{negative semi-definite} & \iff (-1)^k \Delta_k \geq 0 \text{ for all principal minors of order } k = 1, \dots, n. \end{array}$$

**Special case:**  $n = 2$  The quadratic form

$$Q_{\mathbf{A}}(\mathbf{x}) = (x_1 \ x_2) \begin{pmatrix} a_{11} & a_{12} \\ a_{12} & a_{22} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \end{pmatrix} = a_{11}x_1^2 + 2a_{12}x_1x_2 + a_{22}x_2^2$$

- is positive definite if  $a_{11} > 0$  and  $\det \mathbf{A} = a_{11}a_{22} - a_{12}^2 > 0$ ;
- is positive semi-definite if  $a_{11} \geq 0$ ,  $a_{22} \geq 0$  and  $\det \mathbf{A} = a_{11}a_{22} - a_{12}^2 \geq 0$ ;
- is negative definite if  $a_{11} < 0$  and  $\det \mathbf{A} = a_{11}a_{22} - a_{12}^2 > 0$ ;
- is negative semi-definite if  $a_{11} \leq 0$ ,  $a_{22} \leq 0$  and  $\det \mathbf{A} = a_{11}a_{22} - a_{12}^2 \geq 0$ .